



International Guidelines on Vaginal Dilation after Pelvic Radiotherapy

Produced by the International Clinical Guideline Group, chaired
by Dr Tracie Miles, President, National Forum of Gynaecological
Oncology Nurses, UK

International guidelines on vaginal dilation after pelvic radiotherapy

SUMMARY

The purpose of these guidelines is to clarify the evidence and offer expert opinion with respect to the use of vaginal dilation for women having radical or adjuvant pelvic radiotherapy.

This document reviews the literature on the:

- perceived rationale for dilation therapy
- causes and definition of stenosis
- incidence of stenosis
- consequences of stenosis
- prevention of stenosis

Good practice points on the use of vaginal dilation therapy have been developed to guide patients and clinicians. These include:

- Dilation therapy may include the use of dilators, vibrators, fingers, or similar shaped devices. It may not be necessary if vaginal intercourse is resumed weekly (or more) following treatment.
- Dilation therapy should be gentle.
- Dilation therapy may be commenced at approximately 2-8 weeks post radiotherapy, when the acute inflammatory response has settled.
- We suggest that a reasonable duration and frequency of dilation may range from three minutes twice a week, up to ten minutes and twice daily.
- Women may be offered a range of sizes according to their anatomy. It is usual to start with the smallest and progress to whatever size is comfortable.
- Women should know that a small amount of bleeding or 'spotting' after dilator use is normal. If there is a lot of bleeding or pain, a clinician (doctor, nurse or therapy radiographer) should be contacted.
- Review the need for dilation therapy on a regular basis, consider discontinuation of dilation therapy when no longer required; e.g. when sexually active or experiencing no discomfort during vaginal examinations at follow up 1-2 years post treatment.
- If stenosis develops record vaginal toxicity using a recognised score.

International guidelines on vaginal dilation after radiotherapy; a clinical guideline

INDEX

1	Summary
2	Index
4	Clinical guideline group
5	Key to Evidence Statements and Grade of Recommendations
6	Aim and Scope of the Guidelines
6	The Rationale for Vaginal Dilation
7	Cause of Stenosis
7	Defining Vaginal Stenosis
9	Incidence of Vaginal Stenosis
10	The Consequences of Stenosis
11	Literature Review: Use of Dilation to prevent Vaginal Stenosis
13	Interventions to prevent Vaginal Stenosis
15	References

CLINICAL GUIDELINE GROUP

Dr Tracie Miles (Chair), Gynaecology Oncology Clinical Nurse Specialist and University Lecturer, Bristol and Bath, UK.

Membership (in alphabetical order)

- Dr Karin Bergmark, Senior Consultant in Gynaecologic Oncology and in Obstetrics & Gynaecology, Sahlgrenska University Hospital, Gothenburg, Sweden.
- Dr Deborah Watkins Bruner, Robert W. Woodruff Professor of Nursing and Radiation Oncology, Associate Director of Outcomes Research, Winship Cancer Centre, Emory University, United States.
- Sarah Burton, Gynaecology Oncology Clinical Nurse Specialist, Cardiff, Wales.
- Dr Jeanne Carter, Head of Female Sexual Medicine and Womens Health Programme, Memorial Sloan-Kettering Cancer Centre, New York, United States.
- Claire Cohen, Education and Information Co-ordinator, Jo's Cervical Cancer Trust, UK.
- Dr Paul Cornes, Consultant Clinical Oncologist, Bristol, UK.
- Dr Carien L Creutzberg, Consultant Radiation Oncologist, Leiden, Netherlands.
- Liz Darragh, Gynaecology Oncology Clinical Nurse Specialist, Brighton, UK.
- Gail Dunberger, Department of Oncology and Pathology, Karolinska Institutet, Stockholm, Sweden.
- Dr David Edwards, President, British Society for Sexual Medicine, Oxon, UK.
- Emma Elliot, Gynaecology Oncology Clinical Nurse Specialist, London, UK.
- Claire Gildea, Clinical Nurse Specialist in Radiation Oncology. St. Luke's Radiation Oncology Network, St. Luke's Hospital, Dublin, Ireland.
- Pauline Humphrey, Consultant Therapy Radiographer, Bristol, UK.
- Dr Nick Johnson, Pelvic Oncology Surgeon, Bath, UK.
- Lynne Jolicoeur, Advanced Nurse Practitioner, Gynaecology Oncology, Ottawa, Canada.
- Gillian Knowles, Nurse Consultant, Edinburgh Cancer Centre, Scotland.
- Letitia Lancaster, Clinical Nurse Consultant, Gynaecological Oncology, Sydney, Australia.
- Dr Catriona Mclean, Consultant Clinical Oncologist, Edinburgh, Scotland.
- Rosie McCahill, Gynaecology Oncology Clinical Nurse Specialist, Glasgow, Scotland.
- Kathryn Nattress, Royal Prince Alfred Hospital, Sydney Cancer Centre, Australia.
- Lisa Peck, Gynaecology Oncology Clinical Nurse Specialist, Cheltenham, UK.
- Lisa Punt, Consultant Therapy Radiographer, Addenbrookes, UK.
- Professor Roman Rouzier, Obstetrician and Gynaecology Oncology Surgeon, Paris, France.
- Gaby Schwarz, Radioncology & Radiotherapy Department, Universitätsklinikum Heidelberg, Germany.
- Rita Suchy, Radioncology & Radiotherapy Department, Universitätsklinikum Heidelberg, Germany.
- Dr Isabel White, Psychosexual Therapist/Research fellow Cancer Rehabilitation, UK.

Correspondence to the Chair;

Dr Tracie Miles

Department of Gynaecology, Royal United Hospital, Bath, UK

E: traciephd@hotmail.com,

F: +441225825464

T: +1225821553

Page: 7769 via +44 1225428331

FUNDING

No clinical member of the guideline group has any conflict of interest in relation to the development of this guideline. Owen Mumford supported the development of the guidelines by providing meeting rooms, travel expenses for meetings and product expertise.

PREAMBLE

This guideline represents an international consensus on the management of vaginal dilation associated with therapeutic pelvic radiotherapy. The contents are agreed by each member and represent a view based on an interpretation of the available evidence. No reader should assume that this represents the only management option. There are other reasonable care strategies and there is no duty of care for any health care worker to follow these recommendations. Each woman is an individual and holistic care is paramount and there may be many good reasons why these guidelines may not be implemented. The guideline group believes that all women at risk of developing vaginal stenosis as a consequence of therapeutic pelvic radiotherapy are entitled to receive information about the rationale for dilation therapy, and their decision to dilate or not is respected. The guideline group are aware of their responsibility and accept shared responsibility for the publication of these guidelines.

KEY TO EVIDENCE
STATEMENTS
AND GRADE OF
RECOMMENDATIONS

LEVELS OF EVIDENCE	
1++	High quality meta-analyses, systematic reviews of randomised controlled trials (RCTs), or RCTs with a very low risk of bias
1+	Well-conducted meta-analyses, systematic reviews of RCTs, or RCTs with a low risk of bias
1-	Meta-analyses, systematic reviews of RCTs, or RCTs with a high risk of bias
2++	High quality systematic reviews of case control or cohort studies
2+	Well-conducted case control or cohort studies with a low risk of confounding or bias and a moderate probability that the relationship is causal
2-	Case control or cohort studies with a high risk of confounding or bias and a significant risk that the relationship is not causal
3	Non-analytic studies, e.g. case reports, case series
4	Expert opinion

KEY TO EVIDENCE
STATEMENTS
AND GRADE OF
RECOMMENDATIONS
(cont)

GRADES OF RECOMMENDATION	
Note: The grade of recommendation relates to the strength of the evidence on which the recommendation is based. It does not reflect the clinical importance of the recommendation.	
A	At least one meta-analysis, systematic review of RCTs, or RCT rated as 1++ and directly applicable to the target population; Or A body of evidence consisting principally of studies rated as 1+, directly applicable to the target population, and demonstrating overall consistency of results
B	A body of evidence including studies rated as 2++, directly applicable to the target population, and demonstrating overall consistency of results; Or extrapolated evidence from studies rated as 1++ or 1+
C	A body of evidence including studies rated as 2+, directly applicable to the target population and demonstrating overall consistency of results; Or extrapolated evidence from studies rated as 2++
D	Evidence level 3 or 4; Or extrapolated evidence from studies rated as 2+

GOOD PRACTICE POINTS (GPP)

GPP	Recommended best practice based on the clinical experience of the guideline development group
------------	---

AIM AND SCOPE
OF THE GUIDELINES

There are no previous international guidelines concerning the current practice of dilation. Australian practice is variable and tends to be dominated by the non-scientific opinion of the leading clinician¹. North American practice guidance does not recommend dilation during therapy and also advises "patients not to have intercourse during radiation therapy"². The (UK) National Forum of Gynaecological Oncology Nurses (NFGON) collaborated with the European Institute of Health and Medical Sciences to survey UK practice (2004)^{3,10} and recommended routine vaginal dilation for all women receiving pelvic radiotherapy^{4,5}. More flexible advice was issued to UK members in 2010^{6,7}, and scientific uncertainty was emphasised¹⁰.

The purpose of these guidelines is to clarify the evidence and offer expert opinion with respect to the use of vaginal dilation for women having radical or adjuvant pelvic radiotherapy.

THE PERCEIVED RATIONALE FOR VAGINAL DILATION

- Facilitate resumption of sexual relations after radiotherapy;
- Prevent adhesions progression to fibrosis and stenosis of the vagina, especially during the first year after completion of RT (if no intercourse is resumed and the patient is motivated to maintain vaginal patency);
- Allow the medical team to examine and assess the vaginal vault or cervix as part of on-going medical follow up;
- Reduce potential sexual difficulties, e.g. painful sexual intercourse;
- Offer the opportunity to discuss sexual fears/myths associated with pelvic radiotherapy;
- Reduce tissue damage;
- Improve psychological well being.

CAUSE OF STENOSIS

Vaginal stenosis may follow pelvic radiotherapy, especially after the combination of external beam radiotherapy and brachytherapy. Radiation damages vaginal epithelium, connective tissues and small blood vessels causing inflammation and cell death prior to resolution. The subsequent reduced blood supply, tissue hypoxia, loss of elastin, collagen deposition and hyalinisation and fibrosis^{9,10,11} leads to thinning of the vaginal mucosa, loss of lubrication, scarring and fibrosis. This causes a shorter, less elastic and dryer vagina. Some women experience complete loss of a functioning vagina^{12,13}. Additionally, oestrogen deficiency (resulting from radiation induced menopause or natural menopause, or cessation of prior hormone replacement therapy, HRT), may intensify the loss of elasticity and lubrication and thinning and atrophy of the vaginal mucosa¹⁴.

DEFINING VAGINAL STENOSIS

Vaginal stenosis is difficult to define. Nunns et al¹⁵ described vaginal stenosis as the inability to insert two fingers into the vagina. Flay and Matthews¹⁶ defined it as shortening of the vagina to less than 8cm. Bruner¹⁷ et al identified it as a decrease in vaginal length from the normal of 8-9 cm while Schover et al¹⁸ classified it in terms of the vaginal mucosa and vaginal capacity as being 'normal, mildly changed, or severely changed'. Hartman and Diddle¹⁹ graded stenotic changes numerically from one to three. A score of grade 1 represented no stenosis, grade 2 related to stenosis of the upper third of the vagina and grade 3 indicated stenosis of more than the upper third. Greven et al²⁰ mentioned vaginal stenosis in their survey of prognostic factors but did not quantify it as they believed that it was not possible to reliably grade this toxicity. In a systematic review, Denton and Maher²¹ commented on the inconsistency of methods for assessing vaginal changes associated with vaginal stenosis. They also observed wide variability in research rigour among studies reviewed concluding there was no accepted reliable and standard measure of vaginal stenosis.


There is no agreed definition or measurement tool for stenosis; our suggested definition and measurement tools are;

Common Terminology Criteria for Adverse Events (CTCAE), reproductive (and breast disorders) section: vaginal stricture, grade 1 to 3 (grade 3 being vaginal narrowing or shortening interfering with the use of tampons, sexual activity or physical examination).

Late Effects of Normal Tissues, Subjective, Objective, Management, Analytic (LENT SOMA), vagina/sexual dysfunction section.

INCIDENCE OF VAGINAL STENOSIS

The incidence of vaginal stenosis has been reported variably in the literature^{1,22}, ranging from 1.2% to 88%. Authors specifically addressing vaginal stenosis due to brachytherapy report an incidence ranging from 13% to 88%. This is higher than the incidence derived from reports where clinicians simply describe their clinical experience (1.2%-54.7%). This suggests ascertainment bias. The range of data in the scientific and health care literature is illustrated below.



Abitbol & Davenport (1974)²⁴, Abitbol et al. (1996)²⁴, Chadha et al. (1999)²⁵, Decruze et al. (1999)¹², Eltabbakh et al. (1997)²³, Flay & Matthews (1995)¹⁶, Gallion et al. (1987)²⁶, Hart et al. (1997)²⁷, Joslin (1989) (cited in Fieler (1997)²⁸ and extracted from Lancaster, Khor et al. (1985)²⁹, Koumantakis et al. (1998)³⁰, MacLeod et al. (1999)³¹, Nanavati et al. (1993)²⁶, Nori et al. (1994)³³, Noyes et al. (1995)³⁴, Nunns et al. (2000)¹⁵, Ogino et al. (1998)³⁵, Perez et al. (1984)³⁶, Pitkin & Van Voorhis (1971)³⁷, Potter et al. (2000)³⁸, Schover et al (1989)³⁹, Seibel et al (1980)⁴⁰, Souhami et al. (1987)⁴¹, Thomas et al. (1991)⁴², Turner et al. (1998)⁴³, Vasicka et al. (1958), Weiss et al. (1999)⁴⁴.

CERVICAL AND ENDOMETRIAL CANCER

Vaginal shortening after brachytherapy was identified by Bruner et al¹⁷ to be greater for women with cervical than for those with endometrial cancer. It was noted by Flay and Matthews¹⁶ to be more common in those undergoing radiotherapy and surgery for cervical cancer than those undergoing radiotherapy alone. In women who had undergone hysterectomy for endometrial cancer, MacLeod et al³¹ found that the addition of external beam radiotherapy combined with brachytherapy resulted in greater toxicity than the addition of brachytherapy alone. However, Nunns et al¹⁵ found no increased incidence of vaginal stenosis when combining both techniques after hysterectomy. The severity of vaginal stenosis appears to be related to a higher dose per fraction of brachytherapy, an increased number of fractions and a smaller diameter of the brachytherapy applicator⁴⁵.

COLORECTAL/ANAL CANCERS

Hartman and Diddle¹⁹ and Poma⁴⁶ found that neither the age of the patient nor stage of the disease contributed to the degree of vaginal stenosis. Bruner et al¹⁷ observed that women with stage II and III endometrial and cervical cancer experienced increased vaginal shortening compared with those with stage I disease. Pearcey and Petereit⁴⁷ commented that vaginal toxicity is greater when the entire length of the vagina is irradiated rather than the vaginal apex. This observation was supported by Katz et al¹⁰ and Tyree et al⁴⁸ who noted that the lower vagina has a poorer tolerance to radiation than the upper vagina. The effect on the vagina in women who smoke during radiotherapy is unknown but smoking is associated with an increased risk of radiation damage^{49,50}. Brand et al⁶⁴ found that 35% of women in a retrospective case note study had stenosis identified after the 6 month follow up check.

The number of women who develop vaginal stenosis after treatment for anal or low rectal cancer is not well reported. However to help inform and guide the use of recommended dilation therapy it is important to consider the treatment fields in anal and rectal cancer separately.

It is important to note that there has been a changing effect over time of radiation dose, type, technique and addition of chemotherapy. These confounding variables in both assessment and prevalence of stenosis makes comparison with historical studies more difficult.

The addition of chemotherapy to radiotherapy to down-stage locally advanced rectal cancer is now standard practice and may impact long term sexual function⁷⁷.

THE CONSEQUENCES OF STENOSIS

In terms of the impact of vaginal stenosis on quality of life, Nori et al³³ classified vaginal stenosis as a 'minor complication', in the same category as cystitis, proctitis, vaginal necrosis and small bowel obstruction. Abitbol and Davenport²⁴ and Nunns et al¹⁵ identified a relationship between vaginal stenosis and increased levels of discomfort during vaginal examinations. Several authors^{16,17,18,24,25,37,51,52} have postulated an association between vaginal stenosis, the severity of dyspareunia and sexual dysfunction. However, the consequences of stenosis remain individual with some women unaffected by significant damage while others experience long lasting psycho, social and sexual loss from clinically minor damage/vaginal changes.

LITERATURE REVIEW: USE OF DILATION TO MINIMISE VAGINAL STENOSIS

Dilation of the vagina is practiced to prevent or treat vaginal stenosis associated with radiotherapy. Most authors do not make a distinction between treatment or prevention but they should be considered separately.

The current rationale for preventing stenosis is to support a woman who wishes to have penetrative vaginal intercourse and to allow her clinician access to the vaginal vault for clinical surveillance so that early detection of a potentially treatable vaginal recurrence could be detected. The suggestion that dilators will improve quality of survival by easing follow-up, allowing early detection of recurrence and salvage therapy, is not proven.

Textbooks and guidance recommend dilation for the treatment of vaginal stenosis but this does not seem to be supported by conclusive primary research data. For example, Rice⁵², Hassey-Dow⁵³, Faithfull⁵⁴, and Krumm and Lamberti⁵⁵ recommend separating the adhesions between the vaginal walls. Gosselin and Waring¹³ state that "regular use of dilators is essential", Wilmoth and Spinelli⁵⁶ state that vaginal dilation can "almost always" prevent stenosis; Davidson et al⁵⁷ comment on the importance of dilators and Lamberti⁵⁸ "stress that the regular use of dilators usually prevents stenosis".

LITERATURE REVIEW:
USE OF DILATION
TO MINIMISE
VAGINAL STENOSIS
(cont)

Pountney's commentary⁵⁹ and Grigsby et al's¹¹ review both claim that dilators are required to prevent stenosis. Yaniv⁶⁰ expands on this and says that "it stands to reason that we do believe that the use of vaginal dilators, in the right way and time indeed reduce radiation damage such as adhesions". None of these authors offers data in support of their individual recommendations. Burke⁶¹ states that one of the most effective ways to manage vaginal stenosis is the use of dilators and draws on the work of Bransfield et al,⁶². This authority is actually a comparison of dilation use extracted from the case notes of French and American hospitals and contains no data to support the claim. In a previous but now outdated Cochrane review, Denton and Maher²¹ claimed that "use of vaginal dilators to prevent the development of vaginal stenosis is supported by grade IIC evidence". They relied on the case series by Poma of five women treated years after radiotherapy and Decruze's work¹², neither of which address dilation during treatment as a preventative therapy. Brand et al⁷² advocate that dilators should be used within 2 weeks of completion of radiotherapy and also refer to the paper by Decruze. The comments by Abitbol and Davenport²⁴ are more analytical and suggest that "mechanical dilation of the vagina and the use of topical oestrogens appear to be of doubtful value". The Cochrane review by Miles⁶⁵ and her systematic review⁶⁶ concluded that there was no good quality evidence to support dilation therapy during the acute radiation phase but there was level 2 and level 3 evidence that dilation could treat stenosis once it had occurred. A systematic search methodology⁶⁷ found only seven studies with any relevant original data.

Two randomised trials^{68,69}, showed that it was possible to successfully encourage and support women to dilate their vagina but this did not increase sexual function scores. Another trial^{70,71} compared a vibrating dilator with the conventional static one. There were no demonstrable differences in sexual function score, vaginal length or elasticity with either device.

Four observational studies comment on vaginal length and all examine dilation practice after radiotherapy. However, all have limited methodological quality and none have good data on the age of their study population. This limits the conclusions that can be drawn from the results. Decruze et al's report¹² actively promotes a stent that they designed. They showed that their device restored the vaginal length more than could be achieved in historical controls with a traditional device. The assessments were subjective, not blind to treatment allocation and women in the control arm were selected by convenience. Bias is compounded because there is a difference in the comparative groups in age, tumour site and type of radiotherapy but this report suggests that dilation may restore some vaginal length. Poma⁶³ describes five case reports and offers similar evidence. Dilation was used a median of eight years after radiotherapy. These women never had any vaginal penetration following or during their radiotherapy until study recruitment. All five women were able to achieve some vaginal patency after dilation. The study by Sobotkowski et al⁷⁸ was a comparison of 31 women treated with radiotherapy for gynaecological cancer and an unspecified half received Mitomycin C to the top of the vagina with a speculum. Mitomycin C prevents DNA cross linkage and its application via a speculum passage reflects an extreme attempt to separate new adhesions between the vaginal walls. There was no difference in vaginal length with this treatment. Velaskar et al⁷² describes an observational study of vaginal length measured before and after a program of dilation therapy. In this case-series of 89 women, the median length increased from 6 to 10cm over the 4 repeated measures and 46 of 89 women were able to accommodate larger dilators after one year of follow up. It is inappropriate to assume that this is due to dilation

GUIDANCE FOR OFFERING VAGINAL DILATORS

PRINCIPLES OF DILATION THERAPY

HOW TO USE A DILATOR

therapy because it only demonstrates that the vagina recovers length or at least, women are able to tolerate a longer vaginal ruler during clinic visits as the radiation inflammatory changes were subsiding.

There is level 2+ evidence that dilation can be used to treat vaginal shortening once it has occurred but there is no published evidence that it should be used during the acute inflammatory phase in irradiated or recently irradiated mucosa.

Information on dilation therapy should be offered to women to decide if they want to use it. Those who wish to maintain or return to sexual activity after radiotherapy may find dilation therapy supports this. Women who state no need for vaginal patency may prefer not to use dilation therapy. Women may find vaginal examinations at follow up uncomfortable due to the formation of vaginal adhesions. Clinicians may separate the adhesions during examination and may recommend using dilation therapy between follow up examinations to prevent formation of adhesions. This may be appropriate following any pelvic radiotherapy which includes the vagina in the treatment. This may include treatment for cervical, endometrial, vaginal, vulval, anal, low rectal and urological cancers. It is likely that the radiotherapy dose and volume of vagina being treated will impact on the severity of potential vaginal stenosis and selection, where possible, of dilator shape.

The principal of dilation is to expand the vaginal tissues. It is known that stretching healthy skin stimulates mitosis and the development of new epithelial cells. Examples include the plastic surgeons who use tissue expanders with good effect in the breast, scalp and vagina⁷³, and girls born without a vagina (Rokitansky syndrome/or Mayer-Rokitansky-Küster-Hauser Syndrome or müllerian agenesis) can stimulate new skin growth by applying pressure on the müllerian pit (vestibule)^{74,79}. Although the skin of a teenager's vaginal stem cells has no resemblance to epithelial cells after radical radiation, the belief is that the skin will re-grow if stretched, or at least, the vaginal supporting tissues can be encouraged to part if they are stimulated. It follows that there is no science to support one device over another. Any device should suffice. Normal coitus, commercially available sex toys, or clinical dilators should be equally effective. The selection of dilator will depend on choice, local availability and funding arrangements.

GPP Dilation therapy may include the use of dilators, vibrators, fingers, or similar shaped devices.

Instructions for using dilators may differ slightly from hospital to hospital, but the principles are the same. The dilator is inserted gently into the vagina using a lubricant. Once inserted to the top of the vagina to a comfortable point, it is gently rotated and then withdrawn. It is reasonable to assume that dilators must not be applied with force to avoid vaginal/mucosal trauma.

GPP Dilation therapy should be gentle

There is no evidence to support any recommended technique over another. There is no evidence to inform the duration of treatment but some specialists advise use for 6–24 months⁴, or during the first year; others say it should be life-long. Controversies about frequency in the application of

HOW TO USE A DILATOR (cont)

dilation therapy exist. Cartright-Alcarese⁹ advises use of a dilator for 10 minutes a day, 3 times a week. Hartman and Diddle¹⁹ recommended a dildo for women who did not have a “spouse”. Crowther and colleagues⁷⁵ suggested that women should be instructed to apply oestrogen to the vagina (despite the fact radiotherapy destroys hormone receptors) and apply rigorous vaginal dilation and Kegel pelvic floor exercises to promote blood flow to the area. The UK patient charity Macmillan⁷⁶ advise patients that they “can minimise or prevent stenosis by using vaginal dilators from 2–8 weeks after the end of their radiotherapy” and refer to a survey of practice, rather than justifying data⁸. Previous UK guidelines suggested use of dilators for 5 minutes, 3 times a week⁵. There is no evidence to support or refute this. None of these recommendations are supported by data and therefore definitive instructions cannot be issued. However, it seems logical to conclude that the least extreme advice must be more convenient and acceptable for women. Clinicians may discuss acute and late effects of radiotherapy on the vagina and the need for dilators during the consent and planning phase of radiotherapy. Dilation therapy may be introduced during the follow up visits. This could be part of the survivorship program and introduced as an option to support sexual well-being once treatment is over, once the acute local inflammatory phase has started to settle.

GPP Women experiencing difficulty with the concept of dilation/expressing sexual concerns should be offered a comprehensive sexual assessment (where resources available)

GPP Dilation therapy may be commenced at approximately 2-8 weeks post radiotherapy, when the acute inflammatory response has settled.

GPP We suggest that a reasonable duration and frequency of dilation may range from three minutes twice a week, for the first six months, up to ten minutes and twice daily. Once a week thereafter and then occasionally after a year if not experiencing difficulty.

GPP Women may be offered a range of sizes according to their anatomy. It is usual to start with the smallest and progress to whatever size is comfortable. Dilator shape should be determined by the tumour site treated; pointed end for anal, low rectal/ vaginal cancer and flat end for endometrial/ cervix cancer.

GPP Women should be advised that a small amount of bleeding or ‘spotting’ after dilator use is normal. If there is a lot of new bleeding or pain, a clinician should be contacted.

GPP Review the need for dilation therapy on a regular basis, considering discontinuing dilation therapy when no longer required.eg when sexually active or experiencing no discomfort during vaginal examinations at follow up.

GPP If stenosis develops record toxicity using a recognised score.

REFERENCES

- Lancaster L. Preventing vaginal stenosis after brachytherapy for gynaecological cancer: an overview of Australian practices. *European Journal of Oncology Nursing*. 2004. 8(1); 30-39.
- <http://www.cancer.gov/cancertopics/wyntk/uterus/page14>
- White ID, Faithfull S. Vaginal dilation associated with pelvic radiotherapy: A UK survey of current practice. *International Journal of Gynecological Cancer*. 2006. 16(3); 1140-1146.
- Jefferies H, Hoy S, McCahill R, Crichton A. Guidelines on vaginal dilator use after pelvic radiotherapy. *Nursing Times*. 2007. 103(30); 28-29.
- Best Practice guidelines on the use of vaginal dilators in women receiving pelvic radiotherapy. National Forum of Gynaecological Oncology Nurses; <http://www.owenmumford.com/en/range/20/amielle-care.html> last sourced Nov 07.
- Miles T. Round the country update from the president elect. Annual Conference of the National Gynaecological Oncological Nurse Forum. Manchester, July 2010.
- Best practice guidelines for dilator therapy, Autumn news letter. GO news. Published by Owen Mumford, Oxford (in press).
- White I, Faithfull S, Nicholls P. UK Survey of Current Practice in Vaginal Dilatation Associated with Pelvic Radiotherapy. European Institute of Health and Medical Sciences (internal document), 2004. p13.
- Cartwright-Alcares F. Addressing sexual dysfunction following radiation therapy for a gynecological malignancy. *Oncology Nursing Forum* 1995. 22(8); 1227-1232.
- Katz A, Njuguna E, Rakowsky E, Sulkes A, Fenig E. Early development of vaginal shortening during radiation therapy for endometrial or cervical cancer. *International Journal of Gynecological cancer*. 2001. 11(3); 234-235.
- Grigsby PW, Russell A, Bruner D, Eifel P, Koh W-J, Spanos W, Stetz J, Stitt JA, Sullivan J. Late injury of cancer therapy on the female reproductive tract. *International Journal of Radiation Oncology Biology and Physics* 1995. 31(5); 1281-1299.
- Decruze SB, Guthrie D, Magnani R, Prevention of vaginal stenosis in patients following vaginal brachytherapy. *Clinical Oncology* 1999. 11; 46-48.
- Gosselin TK, Waring JS. Nursing management of patients receiving brachytherapy for gynecological malignancies. *Clinical Journal of Oncology Nursing* 2001. 5(2); 59-63.
- Lamb MA. Questions women ask about gynecologic cancer and sexual functioning. *Developments in Supportive Cancer Care*. 1998. 2(1); 11-13.
- Nunns D, Williamson K, Swaney L, Davy M, The morbidity of surgery and adjuvant radiotherapy in the management of endometrial carcinoma. *International Journal of Gynecological Cancer*. 2000. 10; 233-238.
- Flay L, Matthews JHL. The effects of radiotherapy and surgery on the sexual function of women treated for cervical cancer. *International Journal of Radiation Oncology Biology and Physics*. 1995. 31(2); 399-404.
- Bruner DW, Lanciano R, Keegan M, Corn B, Martin E, Hanks GE. Vaginal stenosis and sexual function following intracavitary radiation for the treatment of cervical and endometrial cancer. *International Journal of Radiation Oncology Biology and Physics*. 1993. 27 (4); 825-830.
- Schover LR, Fife M, Gershenson DM. Sexual dysfunction and treatment for early stage cervical cancer. *Cancer*. 1989. 63 (1); 204-212.
- Hartman P, Diddle AW. Vaginal stenosis following irradiation therapy for carcinoma of the cervix uteri. *Cancer*. 1972. 30 (2); 426-429.
- Greven KM, Corn B, Case D, Purser P, Lanciano RM. Which prognostic factors influence the outcome of patients with surgically staged endometrial cancer treated with adjuvant radiation? *International Journal of Radiation Oncology Biology and Physics*. 1997. 39 (2); 413-418.
- Denton A, Maher EJ. Interventions for the physical aspects of sexual dysfunction in women following pelvic radiotherapy (Cochrane Review) 2003. In: The Cochrane Library, Issue 1. Update Software, Oxford. Retrieved 6 February 2003 from: www.cochrane.org/cochrane/revabst/ab003750.html
- Jeffries SA, Robinson JW, Craighead PS, Keats MR. An effective group psychoeducational intervention for improving compliance with vaginal dilation; a randomised trial. *International Journal of Radiation Biology and Physics*. 2006. 6 (4); 404-411.
- Eltabbakh H, Piver MS, Hempling RE, Shin KH. Excellent long term survival and absence of vaginal recurrences in 332 patients with low risk stage I endometrial adenocarcinoma treated with hysterectomy and vaginal brachytherapy without formal staging lymph node sampling: report of a prospective trial. *International Journal of Radiation Oncology Biology and Physics*. 1997. 38 (2); 373-380.
- Abitbol M, Davenport JH. Sexual dysfunction after therapy for cervical carcinoma. *American Journal of Obstetrics and Gynaecology*. 1974. 119(2); 181-189.
- Chadha M, Nanavati PJ, Liu P, Fanning J, Jacobs. 1999. Patterns of failure in endometrial carcinoma stage Ib grade 3 and Ic patients treated with postoperative vaginal vault brachytherapy. *Gynecologic Oncology* 75(1); 103-107.
- Gallion HH, Maruyama Y, van Nagell JR, Donaldson ES, Rowley KC, Yoneda J, Beach JL, Powell DE, Kryscio RJ. Treatment of stage IIIb cervical cancer with Californium-252 fast-neutron brachytherapy and external photon therapy. *Cancer*. 1987. 59 (10); 1709-1712.
- Hart K, Han I, Deppe G, Malviya V, Malone J, Christensen C, Chuba P, Porter A. Postoperative radiation for cervical cancer with pathologic risk factors. *International Journal of Radiation Oncology Biology and Physics*. 1997. 37(4); 833-838.
- Fieler VE. Side effects and quality of life in patients receiving high-dose rate brachytherapy. *Oncology Nursing Forum*. 1997. 24(3); 545-553.
- Khor TH, Chia KB, Chua EJ, Sethi VK, Tan BC, Tan TM. Radiotherapy of carcinoma of the uterine cervix in Singapore, 1973-1975. *International Journal of Radiation Oncology Biology and Physics*. 1985. 11(7); 1313-1316.
- Koumantakis E, Haralambakis I, Koukourakis M, Mazonakis M, Haldeopoulos D, Papageorgiou N, Livas V, Froudarakis G, Varveris H. A pilot study on concurrent platinum chemotherapy and intracavitary brachytherapy for locally advanced cancer of the uterine cervix. *British Journal of Radiology*. 1998. 71 (845); 552-557.
- MacLeod C, Fowler A, Duval P, D'Costa I, Dalrymple C, Elliott P, Atkinson K, Firth I, Carter J. Adjuvant high dose rate brachytherapy with or without external beam radiotherapy post-hysterectomy for endometrial cancer. *International Journal of Gynecological Cancer*. 1999. 9; 247-255.

32. Nanavati PJ, Fanning J, Hilgers RD, Hallstrom J, Crawford D. High dose rate brachytherapy in primary stage I and II vaginal cancer. *Gynecologic Oncology*. 1993. 51(1); 67-71.
33. Nori D, Merimsky O, Batata M, Caputo T. Postoperative high dose rate intravaginal brachytherapy combined with external irradiation for early stage endometrial cancer: a long term follow-up. *International Journal of Radiation Oncology Biology and Physics*. 1994. 30 (4); 831-837.
34. Noyes WR, Bastin K, Edwards SA, Buchler DA, Stitt JA, Thomadsen BR, Fowler JF, Kinsella TJ. Postoperative vaginal cuff irradiation using high dose rate remote afterloading: a phase II clinical protocol. *International Journal of Radiation Oncology Biology and Physics*. 1995. 32 (5); 1439-1443.
35. Ogino I, Kitamura K, Okajima H, Mastubara S. High-dose-rate intracavitary brachytherapy in the management of cervical and vaginal intraepithelial neoplasia. *International Journal of Radiation Oncology Biology and Physics*. 1998. 40(4); 881-887.
36. Perez CA, Breaux S, Bedwinek JM, Madoc-Jones H, Camel HM, Purdy JA, Walz BJ. Radiation therapy alone in the treatment of carcinoma of the uterine cervix. 1984. *Cancer* 54 (2); 235-246.
37. Pitkin RM, VanVoorhis LW. Postirradiation vaginitis: an evaluation of prophylaxis with topical estrogen. *Therapeutic Radiology*. 1971. 99; 417-421.
38. Potter R, Knocke TH, Fellner C, Baldass M, Reinthaller A, Kucera H. Definitive radiotherapy based on HDR brachytherapy with iridium 192 in uterine cervix carcinoma. *Cancer Radiotherapie*. 2000. 4 (2); 159-172.
39. Schover LR, Fife M, Gershenson DM. Sexual dysfunction and treatment for early stage cervical cancer. *Cancer*. 1989. 63 (1); 204-212.
40. Seibel MA, Freeman MG, Graves WL. Carcinoma of the cervix and sexual functioning. *Obstetrics and Gynaecology*. 1980. 55 (4); 484-487.
41. Souhami L, Melo JA, Pareja G. The treatment of stage III carcinoma of the uterine cervix with telecobalt irradiation. *Gynecologic Oncology*. 1987. 28 (3); 262- 267.
42. Thomas H, Pickering DG, Dunn P, Kam KC, Stewart JS, Lambert HE. Treating the vaginal vault in carcinoma of the endometrium using the buchler afterloading system. *British Journal of Radiology*. 1991. 64 (767); 1044-1048.
43. Turner BC, Knisely JP, Kacinski BM, Haffty BG, Gumbs AA, Roberts KB, Frank AH, Peschel RE, Rutherford TJ, Edraki B, Kohorn EI, Chambers SK, Schwartz PE, Wilson LD. Effective treatment of stage I uterine papillary serous carcinoma with high dose rate vaginal apex radiation (192Ir) and chemotherapy. *International Journal of Radiation Oncology Biology and Physics*. 1998. 40 (1); 77-84.
44. Weiss E, Hirnly P, Arnold-Bofinger H, Hess CF, Banberg M. Therapeutic outcome and relation of acute and late side effects in the adjuvant radiotherapy of endometrial carcinoma stage I and II. *Radiotherapy and Oncology*. 1999. 53 (1); 37-44.
45. Sorbe BG, Smeds AC. Postoperative vaginal irradiation with high dose rate afterloading technique in endometrial carcinoma stage I. *International Journal of Radiation Oncology Biology and Physics*. 1990. 18 (2); 305-314.
46. Poma PA. Postirradiation vaginal occlusion: nonoperative management. *International Journal of Gynaecology and Obstetrics*. 1980. 18 (2); 90-92.
47. Pearcey RG, Petereit DG. Post-operative high dose rate brachytherapy in patients with low to intermediate risk endometrial cancer. *Radiotherapy and Oncology*. 2000. 56; 17-22.
48. Tyree WC, Cardenes H, Randall M, Papiez L. High-dose-rate brachytherapy for vaginal cancer: learning from treatment complications. *International Journal of Gynaecological Cancer*. 2002. 12 (1); 27-31.
49. Eifel PJ, Jhingran A, Bodurka DC, Levenback C, Thames H. Correlation of smoking history and other patient characteristics with major complications of pelvic radiation therapy for cervical cancer. *J Clin Oncol* 2002; 20: 3651-3657.
50. Chon BH, Loeffler JS. The Effect of Nonmalignant Systemic Disease on Tolerance to Radiation Therapy. *The Oncologist*, Vol. 7, No. 2, 136-143, April 2002.
51. Bergmark K, Avall-Lundqvist E, Dickman PW, Henningsohn L, Steineck G. Vaginal changes and sexuality in women with a history of cervical cancer. *New England Journal of Medicine*. 1999. 340 (18); 1383-1389.
52. Rice A. Sexuality in cancer and palliative care 1: effects of disease and treatment. *International Journal of Palliative Nursing* 2000; 6(8):392-7.
53. Hassey-Dow K. Altered patterns of sexuality. *Nursing Care in Radiation Oncology*. Philadelphia, WB Saunders, 1992.
54. Faithfull S, Wells M. *Supportive Care in Radiotherapy*. Edinburgh: Churchill Livingstone, 2003.
55. Krumm S, Lamberti J. Changes in sexual behaviour following radiation therapy for cervical cancer. *Journal of Psychosomatic Obstetric Gynaecology* 1993; 14(1):51-63.
56. Wilmoth MC, Spinelli A. Sexual implications of gynecologic cancer treatments. *Journal of Obstetrics and Gynecology and Neonatal Nursing*. 2000. 29(4); 413-21.
57. Davidson SE, Burns M, Routledge J, Swindell R. The impact of radiotherapy for carcinoma of the cervix on sexual function assessed using the LENT SOMA scales. *Radiotherapy and Oncology*. 2003. 68(3); 241-247.
58. Lamberti J. Sexual adjustment after radiation therapy for cervical carcinoma. *Medical aspects of human sexuality*. 1979. 13; 87-8.
59. Pountney D. The loss of femininity, sexual desire and body image that women can suffer after radiotherapy or gynaecological surgery is very real. *Cancer Nursing Practice*. 2005. (4) 9.
60. Yaniv H, Israel Cancer Association ref: 1759, Israel Cancer Association. A new standard in the use of vaginal trainers as a self-care method to minimize pelvic radiation damage.
61. Burke L. Sexual dysfunction following radiotherapy for cancer of the cervix. *British Journal of Nursing*. 1996. 5 (4); 239-244.
62. Bransfield D, Herriot J, Abbotol A. A medical chart for information about sexual functioning in cervical cancer. *Radiotherapy oncology*. 1984. 1 (4); 317-23.
63. Poma PA. Postirradiation vaginal occlusion: nonoperative management. *International Journal of Gynaecology and Obstetrics*. 1980. 18 (2); 90-92.
64. Brand AH, Bull CA, Cakir B. Vaginal stenosis in patients treated with radiotherapy for carcinoma of the cervix. *International Journal of Gynecological Cancer*. 2006. 16 (1); 288-293.

65. Miles T, Johnson N. Vaginal dilator therapy for women receiving pelvic radiotherapy (Protocol). Cochrane Database of Systematic Reviews 2010, Issue 9. Art. No.: CD007291. DOI: 10.1002/14651858.CD007291.pub2.
66. Johnson N, Miles TP, Cornes P. Dilating the vagina to prevent damage from radiotherapy; systematic review of the literature. *BJOG* 2010. 117; 522-531.
67. Higgins JPT and Green S. *Cochrane Handbook for Systematic Reviews of Interventions*. Published by Wiley Cochrane Series. 2008.
68. Jeffries SA, Robinson JW, Craighead PS, Keats MR. An effective group psychoeducational intervention for improving compliance with vaginal dilation; a randomised trial. *International Journal of Radiation Biology and Physics*. 2006. 6 (4); 404-411.
69. Robinson JW, Faris PD, Scott CB. Psychoeducational group increases vaginal dilation for younger women and reduces sexual fears for women of all ages with gynecological carcinoma treated with radiotherapy. *International Journal of Radiation Oncology Biology and Physics*. 1999. 44; 497-506.
70. Miles TP and Johnson N. Systematic literature review, a randomised trial and a preference trial challenging the practice of vaginal dilation during radiotherapy. 13th Biennial Meeting of the International Gynaecological Cancer Society (IGCS 2010), Prague, Czech Republic, October 23-26, 2010.
71. Miles TP and Johnson N. Systematic literature review, a randomised trial and a preference trial challenging the practice of vaginal dilation during radiotherapy. *International journal of Gynaecological cancer*, 2010, in press.
72. Velaskar SM, Martha R, Mahantashetty U, Badakare JS, Shrivastava SK. Use of indigenous vaginal dilator in radiation induced vaginal stenosis. *The Indian Journal of Occupational Therapy*. 2007. 34(1); 3-6.
73. Johnson N, Batchelor A, Lilford RJ. Experience with tissue expansion vaginoplasty. *British Journal of Obstetrics and Gynaecology*. 1991. 98; 564-8.
74. Johnson N, Lilford RJ. The surgical treatment of gynaecological congenital malformations. In: *Progress in Obstetrics and Gynaecology* 8. Studd J edition. Vol. Chapter 25. Edinburgh : Churchill Livingstone, 1990; 413-431.
75. Crowther ME, Corney RH, Shepherd JH. Psychosexual implications of gynaecological cancer. *BMJ*. 1994. 308(6933); 869-70.
76. http://www.macmillan.org.uk/Cancerinformation/Cancertreatment/Treatmenttypes/Radiotherapy/Pelvicradiotherapyinwomen/Sexualeffects.aspx#DynamicJumpMenuManager_6_Anchor_4
77. Das et al (2010) Long-term quality of life after radiotherapy for the treatment of anal cancer. *Cancer* Feb 15; 822-829
78. Sobotkowski J, Markowska J, Fijuth J, Pietraszek A. Preliminary results of mitomycin C local application as post-treatment prevention of vaginal radiation-induced morbidity in women with cervical cancer. *European Journal of Gynaecological Oncology*. 2006. 27(4); 356-8.
79. Morcel, K, Paniel, B-J, Rouizer, R. Sexual and functional results after nonsurgical or surgical creation of a neovagina in women with Mayer-Rokitansky-Küster-Hauser syndrome: a comparative study.

Published by



Owen Mumford Ltd, Brook Hill, Woodstock, Oxon OX20 1TU

June 2012